

Peer-Reviewed Literature

- Bagarinao, T. (1992). Sulfide as an environmental factor and toxicant: tolerance and adaptations in aquatic organisms. *Aquatic Toxicology*, 24(1-2), 21-62.
- Bagarinao, T., & Vetter, R. D. (1989). Sulfide tolerance and detoxification in shallow-water marine fishes. *Marine Biology*, 103(3), 291-302.
- Bagarinao, T., & Vetter, R. D. (1990). Oxidative detoxification of sulfide by mitochondria of the California killifish *Fundulus parvipinnis* and the speckled sanddab *Citharichthys sitemaeus*. *Journal of Comparative Physiology B: Biochemical, Systemic, and Environmental Physiology*, 160(5), 519-527.
- Bill, M., Rhew, R. C., Weiss, R. F., & Goldstein, A. H. (2002). Carbon isotope ratios of methyl bromide and methyl chloride emitted from a coastal salt marsh. *Geophysical research letters*, 29(4).
- Bouma, A. H. (1963). A graphic presentation of the facies model of salt marsh deposits. *Sedimentology*, 2(2), 122-129.
- Bradshaw, J. S. (1968). Environmental parameters and marsh foraminifera. *Limnology and Oceanography*, 13(1), 26-38.
- Bresnahan, P. J., Wirth, T., Martz, T. R., Andersson, A. J., Cyronak, T., D'Angelo, S., ... & Statom, N. (2016). A sensor package for mapping pH and oxygen from mobile platforms. *Methods in Oceanography*, 17, 1-13.
- Cheng, B. S., & Hovel, K. A. (2010). Biotic resistance to invasion along an estuarine gradient. *Oecologia*, 164(4), 1049-1059.
- Crooks, J. A. (2001). Assessing invader roles within changing ecosystems: historical and experimental perspectives on an exotic mussel in an urbanized lagoon. *Biological Invasions*, 3(1), 23-36.
- Crooks, J. A., & Khim, H. S. (1999). Architectural vs. biological effects of a habitat-altering, exotic mussel, *Musculista senhousia*. *Journal of Experimental Marine Biology and Ecology*, 240(1), 53-75.
- Currin, C. A., Levin, L. A., Talley, T. S., Michener, R., & Talley, D. (2011). The role of cyanobacteria in Southern California salt marsh food webs. *Marine Ecology*, 32(3), 346-363.
- Cyronak, T., Andersson, A. J., D'Angelo, S., Bresnahan, P., Davidson, C., Griffin, A., ... & White, M. (2018). Short-Term Spatial and Temporal Carbonate Chemistry Variability in Two Contrasting Seagrass Meadows: Implications for pH Buffering Capacities. *Estuaries and Coasts*, 41(5), 1282-1296.
- DiBacco, C., & Levin, L. A. (2000). Development and application of elemental fingerprinting to track the dispersal of marine invertebrate larvae. *Limnology and Oceanography*, 45(4), 871-880.
- Fleminger, A., & Kramer, S. H. (1988). Recent introduction of an Asian estuarine copepod, *Pseudodiaptomus marinus* (Copepoda: Calanoida), into southern California embayments. *Marine Biology*, 98(4), 535-541.

- Ford, L. D., & Norris, K. S. (1988). The University of California Natural Reserve System: Progress and prospects. *BioScience*, 38(7), 463-470.
- Grover, C. D., Dayton, K. C., Menke, S. B., & Holway, D. A. (2008). Effects of aphids on foliar foraging by Argentine ants and the resulting effects on other arthropods. *Ecological Entomology*, 33(1), 101-106.
- Hentschel, B. T. (2004). Sediment resuspension and boundary layer flow dramatically increase the growth rates of interface-feeding spionid polychaetes. *Journal of marine systems*, 49(1), 209-224.
- Hentschel, B. T., & Herrick, B. S. (2005). Growth rates of interface-feeding spionid polychaetes in simulated tidal currents. *Journal of Marine Research*, 63(5), 983-999.
- Janousek, C. N. (2009). Taxonomic composition and diversity of microphytobenthos in southern California marine wetland habitats. *Wetlands*, 29(1), 163-175.
- Janousek, C. N., Currin, C. A., & Levin, L. A. (2007). Succession of microphytobenthos in a restored coastal wetland. *Estuaries and Coasts*, 30(2), 265-276.
- Johnson, M. R., Williams, S. L., Lieberman, C. H., & Solbak, A. (2003). Changes in the abundance of the seagrasses *Zostera marina* L.(eelgrass) and *Ruppia maritima* L.(widgeongrass) in San Diego, California, following and El Niño Event. *Estuaries and Coasts*, 26(1), 106-115.
- Kelly, C. K. (1990). Plant foraging: a marginal value model and coiling response in *Cuscuta subinclusa*. *Ecology*, 71(5), 1916-1925.
- Krug, P. J. (2001). Bet-hedging dispersal strategy of a specialist marine herbivore: a settlement dimorphism among sibling larvae of *Alderia modesta*. *Marine Ecology Progress Series*, 213, 177-192.
- Krug, P. J., & Zimmer, R. K. (2000). Larval settlement: chemical markers for tracing production, transport, and distribution of a waterborne cue. *Marine Ecology Progress Series*, 207, 283-296.
- Krug, P. J., & Zimmer, R. K. (2004). Developmental dimorphism: consequences for larval behavior and dispersal potential in a marine gastropod. *The Biological Bulletin*, 207(3), 233-246.
- Krug, P. J., Gordon, D., & Romero, M. R. (2012). Seasonal polyphenism in larval type: rearing environment influences the development mode expressed by adults in the sea slug *Alderia willowi*. *Integrative and Comparative Biology*, 52(1), 161-172.
- Levin, L. A. (1980). Life history and dispersal patterns of four infaunal polychaetes: implications for the structure of a dense assemblage. *American Zoologist*, 20(4), 881.
- Levin, L. A. (1981). Dispersion, feeding behavior and competition in two spionid polychaetes. *Journal of Marine Research*, 39(1), 99-117.
- Levin, L. A. (1982). Interference interactions among tube-dwelling polychaetes in a dense infaunal assemblage. *Journal of Experimental Marine Biology and Ecology*, 65(2), 107-119.
- Levin, L. A. (1983). Drift tube studies of bay-ocean water exchange and implications for larval dispersal. *Estuaries*, 6(4), 364-371.

- Levin, L. A. (1984). Life history and dispersal patterns in a dense infaunal polychaete assemblage: community structure and response to disturbance. *Ecology*, 65(4), 1185-1200.
- Levin, L. A. (1984). Multiple patterns of development in *Streblospio benedicti* Webster (Spionidae) from three coasts of North America. *The Biological Bulletin*, 166(3), 494-508.
- Levin, L. A., Talley, T. S., & Hewitt, J. (1998). Macrobenthos of *Spartina foliosa* (Pacific cordgrass) salt marshes in southern California: community structure and comparison to a Pacific mudflat and a *Spartina alterniflora* (Atlantic smooth cordgrass) marsh. *Estuaries and Coasts*, 21(1), 129-144.
- Levin, L. A., & Talley, T. S. (2002). Influences of vegetation and abiotic environmental factors on salt marsh invertebrates. In *Concepts and controversies in tidal marsh ecology* (pp. 661-707). Springer Netherlands.
- Levin, L. A., & Talley, T. S. (2002). Natural and manipulated sources of heterogeneity controlling early faunal development of a salt marsh. *Ecological Applications*, 12(6), 1785-1802.
- Lorda, J., Hechinger, R. F., Cooper, S. D., Kuris, A. M., & Lafferty, K. D. (2016). Intraguild predation by shore crabs affects mortality, behavior, growth, and densities of California horn snails. *Ecosphere*, 7(5).
- Loreau, M. (1998). Biodiversity and ecosystem functioning: a mechanistic model. *Proceedings of the National Academy of Sciences*, 95(10), 5632-5636.
- Macdonald, K. B. (1969). Quantitative studies of salt marsh mollusc faunas from the North American Pacific coast. *Ecological Monographs*, 39(1), 33-60.
- Mahmoudi, N., Modanu, M., Brandt, Y., & Andrade, M. C. (2008). Subtle pedipalp dimorphism: a reliable method for sexing juvenile spiders. *Journal of Arachnology*, 36(3), 513-517.
- Moran, R. (1980). Did you say mangroves? In Mission Bay? *Environment Southwest*, 488(1-4), 1-4.
- Moseman, S. M. (2007). Opposite diel patterns of nitrogen fixation associated with salt marsh plant species (*Spartina foliosa* and *Salicornia virginica*) in southern California. *Marine Ecology*, 28(2), 276-287.
- Moseman, S. M., Zhang, R., Qian, P. Y., & Levin, L. A. (2009). Diversity and functional responses of nitrogen-fixing microbes to three wetland invasions. *Biological Invasions*, 11(2), 225-239.
- Moseman-Valtierra, S. M., Johnson, R., Zhang, R., & Qian, P. Y. (2009). Differences in cordgrass structure between a mature and developing marsh reflect distinct N₂-fixing communities. *Wetlands*, 29(3), 919-930.
- Moseman-Valtierra, S. M., Armaiz-Nolla, K., & Levin, L. A. (2010). Wetland response to sedimentation and nitrogen loading: diversification and inhibition of nitrogen-fixing microbes. *Ecological Applications*, 20(6), 1556-1568.
- Noto, A. E., & Shurin, J. B. (2016). Population variation affects interactions between two California salt marsh plant species more than precipitation. *Oecologia*, 180(2), 499-506.

- Noto, A. E., & Shurin, J. B. (2017). Early Stages of Sea-Level Rise Lead to Decreased Salt Marsh Plant Diversity through Stronger Competition in Mediterranean-Climate Marshes. *PloS one*, *12*(1), e0169056.
- Noto, A. E., & Shurin, J. B. (2017). Interactions among salt marsh plants vary geographically but not latitudinally along the California coast. *Ecology and Evolution*.
- Novoa, A., Talley, T. S., Talley, D. M., Crooks, J. A., & Reyns, N. B. (2016). Spatial and temporal examination of bivalve communities in several estuaries of Southern California and Northern Baja California, MX. *PloS one*, *11*(2), e0148220.
- Phleger, F. B. (1970). Foraminiferal populations and marine marsh processes. *Limnology and Oceanography*, *15*(4), 522-534.
- Phleger, F. B., & Bradshaw, J. S. (1966). Sedimentary environments in a marine marsh. *Science*, *154*(3756), 1551-1553.
- Reusch, T. B., & Williams, S. L. (1999). Macrophyte canopy structure and the success of an invasive marine bivalve. *Oikos*, 398-416.
- Rhew, R. C., Miller, B. R., & Weiss, R. F. (2000). Natural methyl bromide and methyl chloride emissions from coastal salt marshes. *Nature*, *403*(6767), 292.
- Roe, K. L., Hogle, S. L., & Barbeau, K. A. (2013). Utilization of heme as an iron source by marine Alphaproteobacteria in the Roseobacter clade. *Applied and environmental microbiology*, *79*(18), 5753-5762.
- Romero, M. R., Phuong, M. A., Bishop, C., & Krug, P. J. (2013). Nitric oxide signaling differentially affects habitat choice by two larval morphs of the sea slug *Alderia willowi*: mechanistic insight into evolutionary transitions in dispersal strategies. *Journal of Experimental Biology*, *216*(6), 1114-1125.
- Seale, E. M., & Zacherl, D. C. (2009). Seasonal settlement of *Olympia* oyster larvae, *Ostrea lurida* Carpenter 1864 and its relationship to seawater temperature in two southern California estuaries. *Journal of Shellfish Research*, *28*(1), 113-120.
- Siegel, P. R., & Wenner, A. M. (1984). Abnormal reproduction of the sand crab *Emerita analoga* in the vicinity of a nuclear generating station in Southern California. *Marine Biology*, *80*(3), 341-345.
- Smolensky, N., Romero, M. R., & Krug, P. J. (2009). Evidence for costs of mating and self-fertilization in a simultaneous hermaphrodite with hypodermic insemination, the opisthobranch *Alderia willowi*. *The Biological Bulletin*, *216*(2), 188-199.
- Sobecky, P. A., Mincer, T. J., Chang, M. C., & Helinski, D. R. (1997). Plasmids isolated from marine sediment microbial communities contain replication and incompatibility regions unrelated to those of known plasmid groups. *Applied and environmental microbiology*, *63*(3), 888-895.
- Talley, T. S., & Levin, L. A. (1999). Macrofaunal succession and community structure in Salicornia marshes of southern California. *Estuarine, Coastal and Shelf Science*, *49*(5), 713-731.

- Talley, T. S., Crooks, J. A., & Levin, L. A. (2001). Habitat utilization and alteration by the invasive burrowing isopod, *Sphaeroma quoyanum*, in California salt marshes. *Marine Biology*, 138(3), 561-573.
- Talley, D. M., Talley, T. S., & Blanco, A. (2015). Insights into the establishment of the Manila clam on a tidal flat at the southern end of an introduced range in southern California, USA. *PloS one*, 10(3), e0118891.
- Trnka, S., & Zedler, J. B. (2000). Site conditions, not parental phenotype, determine the height of *Spartina foliosa*. *Estuaries*, 23(4), 572-582.
- Weinersmith, K. L., Brown, C. E., Clingen, K. B., Jacobsen, M. C., Topper, L. B., & Hechinger, R. F. (2018). *Euhaplorchis californiensis* Cercariae Exhibit Positive Phototaxis and Negative Geotaxis. *Journal of Parasitology*.
- Whitcraft, C. R., & Levin, L. A. (2007). Regulation of benthic algal and animal communities by salt marsh plants: impact of shading. *Ecology*, 88(4), 904-917.
- Zedler, J. B. (1988). Salt marsh restoration: lessons from California. *Rehabilitating damaged ecosystems*, 1, 123-138.
- Zemal, R., & Fancher, J. M. (1988). Foraging behavior and foods of the light-footed clapper rail. *The Condor*, 90(4), 959-962.

Theses

- Angeloni, L. M. (2001). *Mating strategies of simultaneously hermaphroditic Opisthobranchs: sexual selection, body size and sex allocation* (Doctoral dissertation).
- Boyer, K. E. (1994). *Scale insect damage in constructed salt marshes: Nitrogen and other factors* (Master's thesis).
- Brothers, S. B. (1975). *The comparative ecology and behavior of three sympatric California gobies* (Doctoral dissertation).
- Crooks, J. A. (1998). *The effects of the introduced mussel, *Musculista senhousia*, and other anthropogenic agents on benthic ecosystems of Mission Bay, San Diego* (Doctoral dissertation).
- Hoyt, K. (2009). *Levels of metals from salt marsh plants from Southern California, USA* (Master's thesis).
- Janousek, C. (2005). *Functional diversity and composition of microalgae and photosynthetic bacteria in marine wetlands: Spatial variation, succession, and influence on productivity* (Doctoral dissertation).
- Krug, P. J. (1998). *Chemical and larval ecology of Opisthobranch molluscs: Variable development modes and settlement cues for larvae of *Alderia modesta** (Doctoral dissertation).

Lemons, E. P. T. (2017). *Sediment Accretion Rates and Metal Contamination of the Kendall-Frost Marsh Reserve in Mission Bay, CA* (Master's thesis).

Levin, L. A. (1982). *The roles of life history, dispersal, and interference competition in the population and community structure of a dense infaunal polychaete assemblage* (Doctoral dissertation).

Lipski, D. M. (2005). *Larval supply, settlement and recruitment of fishes in seagrass beds in a Southern California embayment* (Master's thesis).

Macdonald, K. B. (1969). *Quantitative studies of salt marsh mollusk faunas from the North American Pacific Coast* (Doctoral dissertation).

McCray, A. T. (2001). *Halophyte responses to soil variation and interspecific competition in Southern California salt marshes* (Doctoral dissertation).

Phleger, C. F. (1964). *Effect of varying salinity on lipid, protein, and mineral content of a coastal salt marsh plant, *Spartina leiantha** (Doctoral dissertation).

Talley, D. M. (2000). *The role of resident fishes in linking habitats of a Southern California salt marsh* (Doctoral dissertation).

Reports (Governmental, Technical, Student, Etc.)

Bradshaw, J. S. (1968). *A multi-factor instrument for recording ecological parameters* (No. 47). Marine Foraminifera Laboratory, Scripps Institution of Oceanography, University of California, San Diego.

Chaine, A. (1996) *Sexual Selection in *Navanax inermis* a Hermaphroditic Sea Slug*.

Daniel, T. (2016). *Avicennia (Acanthaceae: Avicennioideae) in North America and Mesoamerica. *Proceedings of the California Academy of Sciences*, 63(5), 4th ser., 163-189.*

Deheyn, D. (2009). *Metals concentration in salt marshes plants and kelp around San Diego: A window to environment quality. *UCSD Sustainability Solutions Institute*.*

Dexter, D. M. (1983). *Soft bottom infaunal communities in Mission Bay. *California Fish and Game*, 69(1), 5-17.*

Doherty, D. (1984) *Population density and migration in *Cerithidea californica**.

Dworak, B. (1990) *Urban Runoff: The problem and solutions*.

Janousek, C. N. (2011). *Common benthic algae and cyanobacteria in southern California tidal wetlands. *Scripps Institution of Oceanography*.*

Levin, L. A. (2002). *Alteration of Wetland Habitat by Two Exotic Invertebrates. *California Sea Grant College Program*.*

Levin, L. A., Talley, T. S., Larson, A. A., & Jones, A. (1997). Faunal composition in the Tijuana River estuarine intertidal habitats and the role of life histories in faunal recovery of southern California restored wetlands. *Final report for the National Estuarine Research Reserve System, NOAA Award No. NA 670R0237.*

Phleger, F. B. (1967). Marsh foraminiferal patterns, Pacific coast of North America. *Anales de Instituto Biología Universidad Nacional Autónoma de México*, 38, 11-38.

Reiter, M. E., Shuford, W. D., Strum, K. M., Hickey, C. M., & Page, G. W. (2012). A MONITORING PLAN FOR WINTERING SHOREBIRDS IN COASTAL CALIFORNIA AND NORTHERN BAJA CALIFORNIA, MEXICO Version 1.0.

Rosenfeld, B. (1990). *Crabs at the Kendall-Frost Salt Marsh.*

Saladin, M. (1998). *The influence of the non-native mussel, Musculista senhousia, on sediment properties in Mission Bay, San Diego, CA, USA.*

Sigala, M. (1995) *Competition of Spartina foliosa with succulents in a salt marsh community.*

Stoms, D. M. (2014). Ecological research reserve planning. In *Making Transparent Environmental Management Decisions* (pp. 175-203). Springer Berlin Heidelberg.

Tuckwell, J. (1990). *Proposal for the management, preservation and expansion of the Kendall-Frost Mission Bay Marsh Natural Reserve.*

Varnado, B. (1983) *Distribution of parasite Cuscuta salina on host plant species in a Southern California salt marsh environment.*

Wenner, A. M., Page, H.M., and Siegel, P.R. (1985) Variation in size at onset of egg production. *Factors in adult growth, Crustacean issues 3.* Balkema, Rotterdam, 149–164.

Zemba, R., Hoffman, S. M., & Konecny, J. (2012). Status and Distribution of the Light-footed Clapper Rail in California, 2012 Season. *California Department of Fish and Game, Nongame Wildlife Program Report, 2.*

Zemba, R., Hoffman, S. M., Gailband, C., & Konecny, J. (2016). Light-footed Ridgway's (Clapper) Rail Management, Study, and Zoological Breeding in California 2016 Season.